

SCIENTIFIC DEVELOPMENT AND ACHIEVEMENTS

volume 4

London 2018

Scientific development and achievements

Copyright © 2018 by Hamaiunova V., Hlushko T., Honenko L., Prylipko T., Bukalova N., Bogatko N., Shtuiak A., Dobrovolsky V., Stankevych S., Zabrodina I., Yevtushenko M., Biletskyj Y., Sychov M., Holubiev M., Voloshin N., Golovan L., Klymenko I., Chupryna Y., Lisovyy M., Fedorchuk S., Klymenko T., Trembitska O., Vasilyeva Ju., Lezhenina I., Kochmarskii V., Melnyk V., Skiba M., Vorobyova V., Vereschak V., Pasenko O., Bykov I., Vishnikin A., Al-Shwaiyat Mohammed E.A., Okovytyy S., Chernavskaya A., Hedjazi M., Zhuk T., Vorobyova V., Margarita S., Zabarna I., Prosaniiy S.

ALL RIGHTS RESERVED
Published by Sciemcee Publishing.
LP22772, 20-22 Wenlock Road
London, United Kingdom
N1 7GU

Sciemcee Publishing is part of SCIEMCEE.

It furthers the SCIEMCEE's mission by disseminating knowledge in the pursuit of education, learning and research at the highest international levels of excellence.

No part of this publication may be reproduced in any manner without the express written consent of the publisher, except in the case of brief excerpts in critical reviews or articles. All inquiries be address to Sciemcee Publishing, LP22772, 20-22 Wenlock Road, London, N1 7GU or publishing@sciemcee.com.

First Edition: 2018

A catalogue record for this publication is available from British Library.

Sciemcee Publishing has no responsibility for the persistence or accuracy of URLs for external or third-party internet referred in this publication, and does not guarantee that any content on such websites is, or will remain, accurate or appropriate.

Every effort has been made in preparing this book to provide accurate and up-to-date information which is in accord with accepted standards and practice at the time of publication. Nevertheless, the authors, editors and publishers can make no warranties that the information contained herein is totally free from error. The authors, editors and publishers therefore disclaim all liability for direct or consequential damages resulting from the use of material contained in this book. Readers are strongly advised to pay careful attention to information provided by the book.

Sciemcee Publishing also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

Sciemcee Publishing books may be purchased for educational, business, or sales promotional use. For information, please e-mail the Sciemcee Publishing at publishing@sciemcee.com.

ISBN 978-1-9993071-0-3
Includes bibliographical references and index.



CONTENTS

| | |
|---|----|
| INTRODUCTION | 6 |
| AGRICULTURAL SCIENCES | 13 |
| Hamaiunova V., Hlushko T., Honenko L. PRESERVATION OF SOIL FERTILITY AS A BASIS FOR IMPROVING THE EFFICIENCY OF MANAGEMENT IN THE SOUTHERN STEPPE OF UKRAINE..... | 13 |
| Conclusion..... | 25 |
| Bibliographical references..... | 26 |
| Prylipko T., Bukalova N., Bogatko N. DEVELOPMENT OF PRACTICAL MEASURES AND WAYS OF THEIR REALIZATION FOR CONTROL, MANAGEMENT OF DAIRY RAW MATERIALS AND DAIRY PRODUCTS IN ACCORDANCE WITH EU NORMS | 28 |
| Conclusion..... | 38 |
| Bibliographical references..... | 38 |
| Prylipko T., Shtuiak A., Dobrovolsky V. PHYSICO-CHEMICAL INDICATORS OF SLAUGHTER PRODUCTS FOR THE USE OF VARIOUS SELENIUM SUPPLEMENTS IN THE DIET OF PIGS..... | 41 |
| Conclusion..... | 48 |
| Bibliographical references..... | 49 |
| Stankevych S., Zabrodina I., Yevtushenko M., Biletskyj Y. THE HISTORY OF AGRICULTURAL ENTOMOLOGY DEVELOPMENT IN UKRAINE IN THE XIX-XXI CENTURIES AND THE ROLE OF THE ENTOMOLOGICAL SCHOOL OF KHARKIV NATIONAL AGRARIAN UNIVERSITY NAMED AFTER V.V. DOKUCHAIEV..... | 50 |
| Conclusion..... | 66 |
| Bibliographical references..... | 67 |
| Sychov M., Holubiev M. RATIONING OF MINERAL NUTRITION OF CHICKEN BROILERS..... | 69 |
| Conclusion..... | 78 |
| Bibliographical references..... | 79 |

Voloshin N.

| | |
|---|----|
| OPTIMIZATION OF IRRIGATION NORMS AND OPERATIVE MANAGEMENT OF IRRIGATION WITH PAID WATER USE..... | 81 |
| Conclusion..... | 92 |
| Bibliographical references..... | 94 |

BIOLOGICAL SCIENCES

Golovan L., Klymenko I., Chupryna Y.

| | |
|--|-----|
| GENETIC FUND OF THE GENUS <i>VIGNA SAVI</i> AND IDENTIFICATION OF DONOR SOURCES BY MORPHOLOGICAL CHARACTERISTICS AND DURATION OF THE GROWING SEASON..... | 95 |
| Conclusion..... | 108 |
| Bibliographical references..... | 109 |

Lisovyy M., Fedorchuk S., Klymenko T., Trembitska O.

| | |
|--|-----|
| METHODOLOGICAL APPROACHES TO THE STUDY AND DETERMINATION OF THE SPECIES CONDITION OF THE ENTOMOLOGICAL DIVERSITY OF AGRICULTURAL LANDSCAPES OF THE FOREST-STEPPE UKRAINE..... | 111 |
| Conclusion..... | 122 |
| Bibliographical references..... | 122 |

Vasilyeva Ju., Lezhenina I.

| | |
|---|-----|
| TO THE ISSUE OF THE CULTURAL AMARANTH ENTOMOCENOSIS FORMATION..... | 123 |
| Conclusion..... | 133 |
| Bibliographical references..... | 134 |

CHEMICAL SCIENCES

Kochmarskii V., Melnyk V.

| | |
|---|-----|
| FORMATION OF CALCIUM CARBONATE FROM MODEL SOLUTIONS. THEORETICAL AND EXPERIMENTAL INVESTIGATIONS..... | 138 |
| Conclusion..... | 153 |
| Bibliographical references..... | 154 |

Skiba M., Vorobyova V.

| | |
|---|-----|
| GREEN SYNTHESIS OF MONOMETALLIC, BIMETALLIC NANOPARTICLES AND COMPOSITE MATERIALS: PROPERTIES AND APPLICATIONS..... | 156 |
| Conclusion..... | 165 |
| Bibliographical references..... | 166 |

Vereschak V., Pasenko O., Bykov I.

CO-PRECIPIATION METHOD IN NANOSCALE SCANDIA
STABILIZED ZIRCONIA POWDERS TECHNOLOGY FOR CERAMIC
FUEL CELLS..... 168
Conclusion..... 179
Bibliographical references..... 179

**Vishnikin A., Al-Shwaiyat Mohammed E.A., Okovytyy S.,
Chernavskaya A., Hedjazi M.**

ANALYTICAL EFFECTS BASED ON INTERACTION OF ORGANIC
DYES WITH HETEROPOLY ANIONS, ANIONIC METAL COMPLEXES
AND CATIONIC POLYELECTROLYTES..... 181
Conclusion..... 205
Bibliographical references..... 205

Zhuk T., Vorobyova V., Margarita S.

BY-PRODUCTS OF APRICOT PROCESSING AS A SOURCE OF
FUNCTIONAL COMPOUNDS: ANTIOXIDANTS AND INHIBITOR OF
CORROSION..... 209
Conclusion..... 221
Bibliographical references..... 221

VETERINARY SCIENCE

Zabarna I., Prosaniy S.

INFLUENCE OF PHARMAZIN AND TILOTSIKLINVET ON THE
BROILER CHICKENS' PRODUCTIVITY..... 223
Conclusion..... 235
Bibliographical references..... 235

INTRODUCTION

The monograph includes 15 sections, each of which is devoted to a specific problem of national economy's sustainable development.

The first third section "PRESERVATION OF SOIL FERTILITY AS A BASIS FOR IMPROVING THE EFFICIENCY OF MANAGEMENT IN THE SOUTHERN STEPPE OF UKRAINE". Ukraine is known as one of the richest countries in the world for high-quality fertile soils, which are the main means of its agricultural industry. The changes that have taken place in recent decades with the fertility of the main soil types require significant improvement. After all, this is an essential condition for obtaining guaranteed levels of productivity of the main crops and ecological balance in general. A significant role in providing the state with grain, oilseeds, vegetables, melons and other products belongs to the zone of the southern Steppe of Ukraine, which is known as the breadbasket. The article presents the data of changes in the main indicators of soil fertility, measures of their improvement and preservation, modern approaches to increasing the yield of crops in the conditions of climate change and insufficient resource provision, the ways of growth of plant productivity and efficiency of moisture use.

The second section "DEVELOPMENT OF PRACTICAL MEASURES AND WAYS OF THEIR REALIZATION FOR CONTROL, MANAGEMENT OF DAIRY RAW MATERIALS AND DAIRY PRODUCTS IN ACCORDANCE WITH EU NORMS". Compulsory microbiological criteria should be applied only in relation to those products or in those points of the food chain, which cannot be applied to other, more efficient ones. As well as in the cases, when the microbiological criteria is expected to increase the protection degree offered to the consumer. Microorganisms that make up the components of the criteria must be generally recognized, for example pathogens or indicator organisms that concern the specific food products and technological processes. The obtained results in terms of using the improved method of *Listeria monocytogenes*, *Salmonella* and coagulase-positive *Staphylococcus* detection in milk and dairy products were steady and authentic, so these figures can be used in evaluating the safety of milk and dairy products. In addition, it should be noted that the techniques are simple, economical in execution and its results give certain quality indicators.

The third section "PHYSICO-CHEMICAL INDICATORS OF SLAUGHTER PRODUCTS FOR THE USE OF VARIOUS SELENIUM SUPPLEMENTS IN THE DIET OF PIGS". Purpose – to study the effect of feeding selenium inorganic and organic forms on productivity and quality of pigs. To achieve this goal, it was necessary to solve the following problem: to examine the effect of selenite, selenate and sel-Plex diet on performance and quality indicators pork. Level of the selenium in ration before

0,3 mg/kg dry material to account of the selenite and selenate sodium, as well as sowed; sel-plex raises the daily average increases pig relatively with checking analogue on 6,9- 13,1%. The advantage of the organic form of the selenium (sowed; sel-plex) on selenite and selenate sodium on daily average increase forms 6,2 and 8,3%. the Accompaniment inorganic (the selenite and selenate sodium) and organic (sowed; sel-plex) of the forms of the selenium in ration assists the improvement destructive and meat-greasy quality pig – an increase the destructive mass and destructive output, masses flourishes, increasing of the meat specific gravity in touch under simultaneous reduction of the contents of the fat in touch. Herewith the best accumulation in meat specified element is noted beside animal with organic form of the selenium in ration – sowed; sel-plex.

The fourth section “THE HISTORY OF AGRICULTURAL ENTOMOLOGY DEVELOPMENT IN UKRAINE IN THE XIX-XXI CENTURIES AND THE ROLE OF THE ENTOMOLOGICAL SCHOOL OF KHARKIV NATIONAL AGRARIAN UNIVERSITY NAMED AFTER V.V. DOKUCHAIEV”. A historical essay on the development of agricultural entomology in Ukraine since the nineteenth century up to the present days is presented. The most prominent stages and events on the way of the entomology development in the agrarian sector are highlighted. The prominent personalities that influenced the development of the world and national science on crops protection against the pests are mentioned. The role of the Entomological School of Kharkiv National Agrarian University named after V.V. Dokuchaiev as one of the oldest in Ukraine and Europe is defined separately. The main achievements of the outstanding entomologists of Kharkiv National Agrarian University named after V.V. Dokuchaiev such as I.K. Tarnani, V.G. Averin, O.O. Mihulin, B.M. Lytvynov, O.V. Zakharenko, Ye.M. Biletskyi and others are pointed out. The present state of the development of agricultural entomology in Ukraine and in Kharkiv National Agrarian University named after V.V. Dokuchaiev is described.

The fifth section “RATIONING OF MINERAL NUTRITION OF CHICKEN BROILERS”. Based on the results of the analysis of feed materials on the content of micronutrient elements and conducting scientific and economic experiments on chicken broilers, the most effective source of Manganese, Zinc, Iron and Copper in mixed feeds, which contributes to solving the problem of increasing the quality of feeding and intensifying the growth of young farm birds, has been substantiated. The optimum source of essential elements (Mn, Zn, Fe, Cu) for the bodies of chicken broilers through feeds, the level of their incorporation into mixed feeds have been experimentally determined, and their influence on productivity and feed costs per 1 kg of body weight gain have been investigated. The tasks set in the work have been solved experimentally using analytical (analysis of literature sources), zoo-technical (growth indices, feed costs), biochemical (mineral composition of feed materials) and statistical

(processing of the results obtained) methods of research. The use of mixed feeds in feeding chicken broilers which contained glycinate of Manganese, Zinc, Iron and Copper making 75 % of the needs, contributes to increasing their body weight by 2.5 % and increasing the growth rate by 2.4 %. Feed costs per 1 kg of chicken broilers' growth were 1.5 % lower than the young bird index of the control group.

The sixth section "OPTIMIZATION OF IRRIGATION NORMS AND OPERATIVE MANAGEMENT OF IRRIGATION WITH PAID WATER USE". The article presents the method of calculation and optimization of irrigation norms, which is based on simulation modeling of options and graphical optimization in conditions of different prices for water during the years of different water supply. Estimates of additional net profit from irrigation for the main crops are carried out. The method of operative management of irrigation with the use of water-saving irrigation regimes and the use of a multi-layer model of moisture transfer based on theoretical principles of the thermodynamic approach is proposed. The method solves the problems of soil moisture control during irrigation with high accuracy on the basis of the adaptation of the model parameters to the specific conditions of the field. The reduction coefficient of the two-layer and the parameters of the multilayer models of moisture transfer for operational watering planning are checked and identified. A comparison of the calculated and predicted values of soil moisture, which confirms the reliability and high efficiency of the multilayer model of moisture transfer for operational control of irrigation, is carried out.

The seventh section "GENETIC FUND OF THE GENUS VIGNA SAVI AND IDENTIFICATION OF DONOR SOURCES BY MORPHOLOGICAL CHARACTERISTICS AND DURATION OF THE GROWING SEASON". The complex assesment of the genetic fund of the genus Vigna Savi has been investigated and conducted on interspecific, intraspecific and intrapopulation levels. It has been determined that varieties of vigna in terms of morphological variability are more homogeneous than populations. The collectible specimens were marked by an average level of variation in the duration of the growing season and individual phases of the development (0.5-20.7% depending on the type of vigna and genotype). The established asymmetry and excessive variability of the economic characteristics in the populations of the studied species of vigna indicate the peculiarities of the response of the corresponding genotypes to the action of the environmental factors and can serve as a criterion for the adaptability of the species as a whole and individual collection patterns in particular. The significant level of variability of the economic characteristics in the studied species and individual samples of vigna indicates a significant genetic heterogeneity of the study collection and the plasticity of the genotypes to the action of environmental factors. With the use of the Shannon variety index and the Piel equation index, a high level of total phenotypic variability of the vigna collection is

found on the population and species levels ($H' = 0,40-0,67$, $H'_{\max.} = 1,19$). At the same time, the local populations were more heterogeneous than the varieties of vigna.

The eighth section “METHODODOLOGICAL APPROACHES TO THE STUDY AND DETERMINATION OF THE SPECIES CONDITION OF THE ENTOMOLOGICAL DIVERSITY OF AGRICULTURAL LANDSCAPES OF THE FOREST-STEPPE UKRAINE”. Under the existing powerful entomological school, a thorough systematization of the species diversity of insects of Ukraine has not yet been carried out, which makes it difficult to determine the condition of agrarian biodiversity for the environmental justification of the measures envisaged by the UN Convention. In our opinion, an estimation of the entomofauna of agrolandscapes is possible on the example of the representative sample, which is represented by constant and dominant species according to entomological registries created in the past century. In order to optimally summarize representative samples in the process of analytical research, it is suggested to group the known entomofauna of agrolandscapes into basic life forms, each of which requires adequate methods of counting the number due to the peculiarities of the ecology of each grouping. According to the results of the conducted analytical researches, registers of dominant and constant types of insects of the agrolandscapes of the forest-steppe of Ukraine were compiled for life forms of "geophils" (geobionts, herpetobionts) and phytophiles (hortobionts, dendrobionts). The long-term faunal studies of various agrolandscapes have made it possible to determine the presence or absence of certain species in the entomological gatherings and to compare existing species diversity with literary information.

The ninth section “TO THE ISSUE OF THE CULTURAL AMARANTH ENTOMOCENOSIS FORMATION”. The paper deals with the cultural amaranth entomofauna formation in different regions of cultivation. The data on the species composition of amaranth insects in Central and South America, in Africa, in India and in Europe have been analyzed, general patterns of harmful entomofauna formation have been revealed. The research of amaranth entomocomplex in the Eastern Forest-steppe of Ukraine show, that over 200 insect species are associated with this crop. They include phytophagous insects, which are the pests of amaranth: *Lixus subtilis* Boh., *Aphis fabae* Scop., *Lygus pratensis* L., *L. rugulipennis* Popp., *Scotia segetum* Schiff., *Chlorita* sp., *Neoliturus fenestratus* H.-Sch., *Agallia venosa* Four. as well as their entomophags and anthophilous insects. The role of amaranth in increasing the biodiversity of insects in the field crop rotations of Ukraine has been proved.

The tenth section “FORMATION OF CALCIUM CARBONATE FROM MODEL SOLUTIONS. THEORETICAL AND EXPERIMENTAL INVESTIGATIONS”. Kinetics of CaCO_3 crystallization in calcium hydrocarbonate system (CHCS) at $\text{pH} < 10$ is studied. To describe the processes the quasiequilibrium and dynamic models are used. It has been shown that in quasiequilibrium

approximation CHCS is completely described by two parameters - concentration of dissolved CO₂ and pH. Other parameters and their change in time can be calculated by these two parameters. Calculations are compared to data of pCa₂, pCO₂, pCO₃ and pH measurements. It has been shown that transitive processes are incorrectly reflected by quasiequilibrium model. Such calculations are adequate to experimental data of CaCO₃ formation process at degassing of previously saturated with CO₂ system only at final stage of crystallization. In dynamic model kinetic equation of CaCO₃ formation clearly depends on all CHCS parameters but the equation form is determined by pH. Within the range of 6 < pH < 9.5 kinetics of CaCO₃ formation is based on the reaction $\text{Ca}^{2+} + 2\text{HCO}_3^- \rightarrow \text{CaCO}_3\downarrow + \text{HOH} + \text{CO}_2\uparrow$. It has been shown that the process is controlled by diffusion except the time from degassing start to the moment of total crystallization. At final stage of the process metastable CHCS state is formed.

The eleventh section “GREEN SYNTHESIS OF MONOMETALLIC, BIMETALLIC NANOPARTICLES AND COMPOSITE MATERIALS: PROPERTIES AND APPLICATIONS”. In this study we report the potential use plasma-chemical discharge and food waste such as grape skin obtained after grape processing to produce monometallic (AgNPs, AuNPs) and bimetallic nanoparticles (Ag/Au nano-alloy). The nanoparticles were synthesised according to the rules of green chemistry and integrating the goals of green nanotechnology. The synthesized nanoparticles were characterized by UV–Visible spectroscopy, dynamic light scattering. The biosynthesized AgNPs, AuNPs, and Ag-Au nano-alloy showed good antimicrobial activity against Gram-positive and Gram-negative bacteria. AgNPs, AuNPs, Au/Ag bimetallic nanoparticles were formed for heterogeneous and homogeneous composites, respectively. Moreover, the different compositions of as-synthesized Au/Ag bimetallic nanoparticles were utilized for the catalytic reduction of 4-nitrophenol.

The twelfth section “CO-PRECIPIATION METHOD IN NANOSCALE SCANDIA STABILIZED ZIRCONIA POWDERS TECHNOLOGY FOR CERAMIC FUEL CELLS”. This paper deals with physicochemical processes of the formation of nanostructure in stabilized zirconia powders while obtaining it from aqueous solutions of zirconium and scandium salts by coprecipitation method. It was established that precipitation of scandium begins at lower pH value (~ 1) than for individual scandium precipitation that occurs at pH levels ranging from 4.5 to 6. The NMR study showed that in zirconium hydroxides with scandium addition the removal of pore water and structural OH groups is a major obstacle for formation of large crystals and phase transformations.

The thirteenth section “ANALYTICAL EFFECTS BASED ON INTERACTION OF ORGANIC DYES WITH HETEROPOLY ANIONS, ANIONIC METAL COMPLEXES AND CATIONIC POLYELECTROLYTES”. Ion association

complexes (IAs) formed between cationic or anionic organic dyes and large ions originating from inorganic or organic compounds are widely used in analytical chemistry. The extraction is commonly used to separate dye excess that leads to numerous disadvantages. Non-extraction spectrophotometric methods using specific IAs are considered. A theoretical explanation was proposed for the analytical effects observed by the formation of IAs of dyes with heteropoly anions, iodide or rhodanide metal complexes and polyelectrolytes. Most of the found phenomenons are satisfactorily explained in the frame of exciton theory. Approaching of the dye ions can occur due to the formation of precipitate of IA between cationic dyes and heteropoly anions or anionic metal complexes or formation of ion-association bond between anionic dyes and closely placed positively charged groups of cationic polyelectrolytes. Changes in the absorption spectrum are caused then by π - π interaction of aromatic systems of dyes and the formation of J- or H-aggregates.

The fourteenth section “BY-PRODUCTS OF APRICOT PROCESSING AS A SOURCE OF FUNCTIONAL COMPOUNDS: ANTIOXIDANTS AND INHIBITOR OF CORROSION”. Environmentally-friendly tactics become one of the key standard of modern industry. One of them is the practice of employing waste of plants that is present in huge amounts for further industrial applications. Its utilization as promising alternative to environmentally-toxic traditional chemical inorganic corrosion inhibitors (synthetic) received a special attention. Here the protection performance of self-assembled monolayers (SAMs) formed by apricot pomace extract (APE) on the surface of steel has been studied. Characterization of the apricot pomace extract was carried out using Fourier transform infrared spectroscopy (FTIR) and gas chromatography-mass spectrometry (GC-MS) analysis. The antioxidant properties of APE were studied by using a series of methods (total antioxidant activity, reducing power). The anti-corrosion effect of the SAMs was determined by weight loss experiments, scanning electron microscopy, FT-IR and electrochemical methods.

The fifteenth section “INFLUENCE OF PHARMAZIN AND TILTSIKLINVET ON THE BROILER CHICKENS’ PRODUCTIVITY”. Chemicals of the macrolide group – pharmazin and tiltsiklinvet – are widely used in veterinary practice in poultry farming for a treatment or a prevention, in particular in Ukraine. Four groups of broiler chickens of the Cobb 500 day-old cross-breed were formed: two control and two experimental (12 broiler chickens each). Chickens of the first experimental group were fed with pharmazin containing the active substance (AS) tylozine tartrate 500 mg per 1 g, and the second – with a tiltsiklinvet containing the (AS) tylozine tartrate and doxycycline glycate of 100 mg per 1 g of powder. These antibiotics were fed to broiler chickens with preventive purpose for the first 3 days of life, 28–29 and 38–42 days of the experiment. It has been established that the use of pharmazin and tiltsiklinvet positively affects the productivity of chicken broilers. As

a result, one can observe a significant increase in weight of broiler chickens before a slaughter ($p \leq 0.001$, $p \leq 0.01$). Slaughter yield increased by 8,2% when feeding pharmazin, and tilotsiklinvet – by 4.3 %. Also, the weight of all the edible internal organs increased when using pharmazin and tilotsiklinvet ($p \leq 0.05$) and the weight of the liver and muscular part of the stomach when using tilotsiklinvet ($p \leq 0.05$).

Hamaiunova V.

Doctor of agricultural Sciences, Professor, Mykolaiv NAU, Mykolaiv, Ukraine

Hlushko T.

Candidate of agricultural Sciences, Associate Professor, State Higher Educational Institution "Kherson SAU", Kherson, Ukraine

Honenko L.

Candidate of agricultural Sciences, Associate Professor, Mykolaiv NAU, Mykolaiv, Ukraine

PRESERVATION OF SOIL FERTILITY AS A BASIS FOR IMPROVING THE EFFICIENCY OF MANAGEMENT IN THE SOUTHERN STEPPE OF UKRAINE

Introduction. Successful work of the farmer consists primarily in obtaining sustainable productivity of crops and preserving soil fertility – the most important and unique natural resource and the main means of production and the basis of life of all living things on Earth. It is the fertility of the soil and the conditions of management that determine not only the amount of the grown crop, but also the indicators of its quality, the state of the ecological environment. At the same time, in modern conditions of the agricultural sector of Ukraine, there is an urgent need to restore natural ecosystems, maintaining their biological balance at a level that will guarantee the stability of the environment, protection of land from degradation, erosion and loss of fertility.

The area of agricultural lands of Ukraine is one of the largest among the countries of Europe, and their qualitative composition and biological productivity are the richest in the world. We believe that in Ukraine is fertile soils dominate, but unfortunately, recently we have seen their depletion, deterioration of basic properties, loss of productive capacity. These negative changes are mentioned by the researchers of Ukraine. After all, as academician of RAN H.V. Dobrovolskyi noted, the processes of destruction and degradation of the ground cover are non-renewable and determine the development of the ecological situation, which the scientist called a quiet crisis of the planet [1].

Among the problems in the agricultural sector the most important one is to ensure sustainable agriculture, increase its productivity by raising soil fertility and the ability to form a guaranteed crop yield with high quality.

Results. In order to maintain soil fertility in the southern Steppe zone of Ukraine, according to the results of research, 7-8 tons of organic and 80-100 kg of mineral fertilizers should be applied to each hectare of agricultural land and, accordingly, 12-15 tons and 240-260 kg - under irrigation conditions. Such a number of fertilizers in

recent years are not added, and crop production is formed mainly due to the residual reserves in the soil of nutrients and mineralization of humus, so that its content is reduced, the soils gradually get worse, depleted and degraded. This requires an urgent response of agricultural producers, because if it is impossible to restore the spent resources, the degradation processes become irreversible [2,3].

So, modern agricultural use of land requires the development and justification of agricultural and reclamation activities aimed at the preservation and reproduction of soil fertility. This leads to the need for research to predict changes in quantitative and qualitative indicators of natural factors of fertility, patterns of direction and speed of soil processes. It is necessary to effectively and continuously monitor the state of soil fertility, the degree of its erosion, the reaction of the environment, the salt regime of the soil. Recently, such processes as soil contamination with heavy metals, radionuclides, pesticides and other toxicants have also caused concern [4]. Depending on the intensity of land use, including irrigation, soil fertility and the environmental situation in general change.

The most cheap, affordable and cost-effective way of addressing improvement of soil fertility and increasing crop yields is the introduction into production of science-based crop rotations, which due to plaintive-root residues enrich the soil with organic matter, and in case of the selection of legumes it is also free source biological nitrogen. Reasonable alternation of crops improves water and nutrient regimes of the soil, reduces contamination of fields, the presence of pests and pathogens, that is, it contributes to the preservation of ecological balance and rational use of agricultural land [5].

Unfortunately, in recent years, the structure of crop rotations is significantly violated. Thus, during the period 1990-2015 in Ukraine there was a redistribution of acreage under crops, in particular: acreage of forage crops decreased by 4.6 times, which adversely affected the development of forage and livestock industry. During this period, the area under crops of highly profitable export – oriented crops has increased significantly: sunflower – by 2.8 times, rapeseed – by more than ten times, which violates the system of crop rotation and leads to complete depletion of the soil cover and its drying. On the contrary, the acreage under potatoes and vegetables and melons decreased by 5%; now they are concentrated mainly in the private sector.

The main problems of the crop industry, in our opinion, are related to:

- expansion of acreage for cultivation of soil crops-sunflower and rapeseed;
- reduction of crops of fodder group, including perennial grasses, in particular alfalfa, individual cereals (millet, buckwheat), legumes and sugar beet;
- low yield of crops compared to other countries in which their high productivity provides the use of modern innovative and resource-saving technologies;
- concentration of production of fruits and vegetables in households that do not

have appropriate storage facilities and conditions for its storage;

- lack of infrastructure for the sale of such products through fruit and sheep farms;
- high energy and resource costs of crop production;
- lack of innovative approaches to modern methods of tillage and cultivation of crops (systems No-Till, Mini-Till, Strip-Till, organic production, drip irrigation);
- a certain shortage of highly skilled workers who are able to work with the latest means of labor and use high-tech operations for growing crops;
- insufficient return of nutrients to the soil, which does not allow to maintain and improve their condition and ability to form crop yields.

Soil fertility is particularly improved due to inclusion grain legumes in the selection of crops in crop rotations. They have a biological feature to form active complexes with soil microorganisms, which bind a significant amount of nitrogen from the air in the process of nitrogen fixation. It is known that during the growing season legumes, depending on the moisture conditions, accumulate from 60-70 to 150 kg of nitrogen. To meet the needs of plants in this amount of nitrogen is necessary to make 200-500 kg / ha of ammonium nitrate. In addition, symbiotic nitrogen fixation has a much higher economic efficiency than the cost of mineral fertilizers. The nitrogen fixed by legumes is used by plants for several years in full, and is not lost as from mineral nitrogen fertilizers. In the root zone legumes produce an active complex of microorganisms that grow very quickly and divide, resulting in the accumulation of organic matter, which eventually turns into humus substances [5]. Additionally, the root allocation of legumes are capable of dissolving phosphates and increase the assimilation of phosphorus compounds [6]. In general, soil enrichment with organic matter improves plant nutrition with minerals in an accessible form, the soil acquires a dynamic balance.

The research was carried out on dark chestnut soil and southern chernozemic soil which is typical for the South of Ukraine on the fields of the Institute of irrigated agriculture of NAAS and Mykolaiv National Agrarian University according to the generally accepted methods of experimental work.

It was found that in the dark chestnut soil in the area of the Ingulets irrigation system, during four rotations of 7-field rotation with alfalfa, the share of which is 35.7% annually loss of humus from the arable layer of the unfrozen soil is 70-90 kg/ha, and during irrigation they were bigger. In case of complete fertilizer of each rotation crop (in doses that are recommended for the zone) humus content in irrigated soil is stabilized, and through the use of once per rotation and even 80 t/ha of half-decay manure it even grows (Table 1).

The deficit in organic fertilizers in the absence of pus in sufficient quantities in crop rotations should be covered by post-harvest residues and by using by-products of crops for fertilizer. The role of organic fertilizers is extremely important in the

preservation of water-physical properties of soils. Without their use, it is compacted, its water absorption capacity deteriorates, the number of water-resistant units decreases. According to our three years of research non-fertilized dark chestnut soil per hour absorbs 14.72 mm of water at green manure in the year of effect and aftereffect, the proportion rose to 16.3-20.6 per cent, and straw – 22.8-34.6 per cent.

Table 1 Humus content in 0-30 layer of dark chestnut soil under the influence of long-term use of fertilizers in irrigated crop rotation

| Experience option | Humus content, % | | Loss or increase of humus | |
|--|------------------------------|---|---------------------------|-------------------------|
| | for the period of experience | after the 4 th rotation of the crop rotation | in absolute percentage | average annual, kg / ha |
| Without fertilizer | 2.26 | 2.11 | -0.15 | -277.5 |
| P ₂ O ₅ | 2.26 | 2.19 | -0.07 | -129.5 |
| N | 2.26 | 2.23 | -0.03 | -55.5 |
| NPK | 2.26 | 2.25 | -0.01 | -13.5 |
| NPK +80 т/ ha of manure 1 time per rotation of crop rotation | 2.26 | 2.35 | +0.09 | +166.5 |

We have determined that for the incorporation of straw into the soil, even in dry years, moisture accumulates in the arable layer by 15-20% more than without it. This should oblige producers of agricultural enterprises in any case not to burn straw, and to use all post-harvest residues for soil enrichment with organic matter. In addition, each ton of straw put in the soil is equivalent to 4-5 t / ha of manure by the amount of dry organic matter introduced.

Many researchers determined that the amount of by-products, which can be used as fertilizer after harvesting the main crop equals on average to 5-6-10 t/ha. All macro - and micronutrients go with it to the soil. So, wheat straw on 1 hectare produces on average 20 kg of nitrogen, 10 kg of phosphorus and 30-35 kg of potassium, with the remains of corn 45; 18 and 100 kg, and sunflower – 40; 20 and 130 kg respectively. In addition, by-products of agricultural crops contain calcium, magnesium, sulfur and all trace elements. It is also known that the decomposition of crop residues depleted to nitrogen is significantly accelerated by adding, for example, to straw 7-10 kg/t of the active substance of nitrogen or modern biodestructors of stubble [7].

The calculations determined that the use of crop residues and straw, compared with manure, can save on each hectare up to 170 kg of diesel fuel and 15-17% of cash costs.

It is known that without the use of organic and mineral fertilizers to a greater extent irrigated soils, which are formed significantly higher levels of crop yields would deplete and, accordingly, the removal of nutrients will significantly increase. Our

studies have found that the use of organic fertilizers has a positive effect on the main elements of soil fertility in the link of irrigated grain-vegetable crop rotation (corn, winter wheat + post-harvest, tomatoes). Under the influence of straw and green fertilizer, and especially when using both types the content of organic matter, total and water-soluble humus in the soil increases (Table 2).

Table 2 The content of humus and organic matter in the arable layer of dark chestnut soil in the link of irrigated grain-vegetable crop rotation (under tomatoes, average for 3 years)

| Experience option | Term of determination | Content | | |
|--|-----------------------|-------------------|-----------------|--------------------------------|
| | | organic matter, % | common humus, % | water-soluble humus, mg / 100g |
| N ₁₂₀ P ₉₀ K ₆₀ -background | 1 | 6.64 | 2.00 | 17.99 |
| | 2 | 7.21 | 2.04 | 25.81 |
| Background+green fertilizer | 1 | 6.81 | 2.02 | 18.86 |
| | 2 | 7.27 | 2.06 | 26.46 |
| Background + winter wheat straw | 1 | 6.82 | 2.03 | 18.42 |
| | 2 | 7.31 | 2.08 | 25.39 |
| Background+green manure+straw | 1 | 6.91 | 2.04 | 19.02 |
| | 2 | 7.38 | 2.09 | 27.16 |

HIP₀₅ 0,22-0,27 0,01-0,03 1,12-2,47

Note: *) Term of determination: 1 – for the period of sowing-rising; 2 – when harvesting.

The latter, as it is known, characterizes the presence of organic substances in the soil at the primary stages of mineralization. Over time, these organic substances in the process of decomposition will turn directly into humus.

The use of straw of grain crops as an organic fertilizer was also studied on the southern chernozem in various parts of the crop rotation in the cultivation of drought-resistant crop without irrigation (Table 3).

It was established that in the conditions of rain – fed agriculture straw effectively affects the main indicators of soil fertility-the content of humus, organic matter, mobile forms of phosphorus and especially potassium, its structural indicators, microbiological activity and the like. In addition, as it is shown in table 3, the grain yield levels for the combined use of straw and N₆₀P₄₀ are formed above the control, but slightly inferior to mineral nutrition. This is due to the fact that for the decomposition of straw, nitrogen fertilizer addition was not made, and due to increased mineralization of organic substances cellulose-desructive microorganisms that use the nitrogen to decompose organic matter.

Table 3 Influence of fertilizers and predecessors on grain yield of soriz, t/ha


| Crop rotation chain (factor A) | Feed backgrounds (factor B) | | | | | | | | |
|-----------------------------------|-----------------------------|------|------|---|------|------|---------------------------------|------|------|
| | Without fertilizer | | | Straw + N ₆₀ P ₄₀ | | | N ₆₀ P ₄₀ | | |
| | 2004 | 2005 | 2006 | 2004 | 2005 | 2006 | 2004 | 2005 | 2006 |
| pea – barley – soriz | 4.26 | 3.18 | 3.30 | 5,29 | 4,41 | 4,50 | 5,42 | 4,55 | 4,61 |
| sunflower-winter barley-soriz | 4.05 | 2.98 | 3.11 | 5,00 | 4,09 | 4,16 | 5,14 | 4,22 | 4,28 |
| corn- winter barley -soriz | 4.06 | 3.01 | 3.13 | 5,17 | 4,23 | 4,30 | 5,27 | 4,33 | 4,40 |
| winter barley-corn-soriz | 3.95 | 2.88 | 3.02 | 4,92 | 3,93 | 4,12 | 5,12 | 4,16 | 4,24 |
| winter wheat-sunflower-soriz | 3.81 | 2.70 | 2.82 | 4,82 | 3,92 | 4,10 | 5,09 | 4,11 | 4,20 |
| Factor B average | 4.03 | 2.75 | 3.08 | 4,84 | 4,12 | 4,24 | 5,21 | 4,27 | 4,35 |
| HIP ₀₅ , t/ha | 2004 | | | 2005 | | | 2006 | | |
| Factor A | 0.071 | | | 0.051 | | | 0.12 | | |
| Factor B | 0.084 | | | 0.098 | | | 0.19 | | |
| Interaction of factors AB | 0.130 | | | 0.270 | | | 0.21 | | |

It should be noted that organo-mineral nutrition is much better even compared with a mineral background that formed the indicators of quality of soriz grain: it increases the content of protein, starch, total sugars, weight of 1000 grains and the like. At the same time, as determined by us, in the aftereffect, the yield of subsequent crops, which were placed after the cut in all parts of the crop rotation, for the joint use of straw and mineral fertilizers was higher as compared to the control and background of nitrogen-phosphorus fertilizer.

Therefore, the basis of any system of crop production is crop rotation. It is the leading part of the zonal system of agriculture and effective agroecological factor. At the same time, in modern technologies of growing crops, the role of the predecessor is decisive, since elements of the technology of growing the next crop depend on it. Proper crop rotation allows the use of a rational system of fertilizer and soil treatment on the basis of which during rotation to maintain optimal water and nutrient regimes for plants to successfully clear the fields from weeds, pests and diseases, is much more efficient to store and use the moisture, fertilizers and thus improve soil fertility for minor expenses while using performance. We show this by the example of growing winter crops on different predecessors (Fig. 1). These data illustrate the advantages of black steam compared to corn and stubble predecessor.

When applying mineral fertilizers on the natural backgrounds of predecessors in the recommended doses for the Steppe zone of Ukraine, grain yield increases significantly, and more significantly for non-steam more depleted predecessors. Of course, for obtaining the highest levels of grain yield, it is much more effective in this field to use the reserves of accumulated soil moisture and precipitation of the growing season, which is extremely important for the arid zone of the southern Steppe of Ukraine. For the cultivation of crops in modern conditions of management and certain

climate changes, the limiting factor in the formation of their high productivity is the provision of plants with moisture [8].



Notes: 1 – winter wheat (2008-2006 and 2015);
 2 – barley (2014-2016);
 3 – triticale (2014-2016);
 4 – rye (2014-2016);
 5 – average for all grains

Figure 1. The yield of winter crops depending on the predecessor and feed background

Technological methods of growing crops should be directed to the maximum possible accumulation of moisture in the soil and its use by plants for the formation of the crop. The accumulated moisture in the soil for the period of sowing of any crop and precipitation of the vegetation period are used much more productively on fertile soils, for the introduction of organic and mineral fertilizers, that is, for the optimization of plant nutrition.


On depleted soils, among the limiting factors, the second place after moisture is taken by the provision of plants with accessible nutrients and, above all, nitrogen. However, as a result of insufficient amount of fertilizers (especially organic ones) and deterioration of soil fertility there is a need to develop new approaches to their nutrition [9,10].

One of the ways of optimizing plant nutrition for these conditions should become more widely applied to the background of moderate doses of mineral fertilizers,

modern growth-regulating substances for the treatment of seeds before sowing and planting of plants at key periods of the growing season. We have conducted research in this direction and continue to perform it on a number of crops: barley, wheat, triticale, red, soybean, peas, chickpeas, sunflower and other spring and winter crops. The effectiveness of this approach to nutrition optimization is shown by the example of spring wheat (Fig. 2).

These data convince of a significant increase in grain yield in comparison with inconvenient control. Moreover, from application to sowing of N₃₀ P₃₀ it increased on average over the three years from 1.72 to 2.72 g/ha. Maximum yield values are reached for nitrogen fertilization on the background of N₆₀ P₃₀ just in time for planting (3.26 t/ha) and N₃₀ P₃₀ before sowing + N₃₀ at the feeding in the early exit of plants in the tube (3.30 t/ha). Almost at the same level, it was formed by double treatment of plants with D2 (3.08 t/ha) and escort (3.10 t/ha).

With presowing treatment of seeds with an escort, the grain yield was formed even higher, which is convincingly illustrated by Fig. 2.



Notes: phase 1 - exit of plants in up;
phase 2-the beginning of earing

Figure 2 Grain yield of spring wheat depending on the background of the supply and use of regulating substances t/ha (average for 2014-2016)

Studies with growth-regulating drugs were conducted in 2016-2017 with two varieties of spring barley – Stalker and Vakula. In addition to D₂M and Escort bio, fresh florid was used to optimize nutrition in doses of 200 and 300 g / ha and Fresh energy (200 g/ha), which are recommended for spring barley. On average, over the years of research Stalker provided in the control of 2.78 t/ha, Vakula – 2.77 t/ha of grain, with the use of all studied biopreparations on average for all 15 options, the yield increased to 3.59 t/ha and 3.57 t / ha or increased to control without fertilizers by 29.1 and 28.9%, respectively. However, it should be noted that in 2016, the higher grain yield was formed by the Vakula – 4.31 t/ha (the weighted average for all variants of the experience), and in the arid 2017 – only 2.73 t/ha (Table 4). That is, the sort of spring barley Stalker is more plastic and is able to form a stable performance with less dependence on weather and climatic conditions of the year.

Similar results were obtained in the treatment of seeds and plants with bio substances in the cultivation of two varieties of peas Oplot and Tsarevich, the yield of which for three years from these substances increased by an average of 28-30%. At the same time, as in the cultivation of spring crops, the main indicators of the quality of pea grain are significantly improved, the number and mass of nodule nitrogen-fixing bacteria on the roots of plants increases, moisture is effectively used, the cost recovery for the introduced resource-saving measures in the cultivation of this legume crop increases. Similar results were obtained on two varieties of chickpea-Rosanna and Memory.

The same data on the optimization of nutrition through the use of bio substances were obtained for the cultivation of sunflower (grade Dragan). Also in studies conducted in 2016-2017 without fertilizers formed 2.45 t/ha, on the background of the application N₁₅P₁₅K₁₅ (100 kg/ha of NPK) – to 2.94 t/ha, and in the processing plants in the phase of 8-10 leaves and education of the basket, depending on the selection Restigouche substances from of 3.02 to 3.48 t/ha. Under the influence of this measure increases the mass of 1000 seeds, test weight, fat content in grain and the conditional yield of oil per unit area, and alkalinity, on the contrary, decreases.

Thus, as it is known and confirmed by our studies with many crops, to increase their productivity and main indicators of the quality is possible by the application on the background of moderate doses of fertilizers and modern growth-regulating substances. This measure provides nutrition of young plants with macro-, microelements and partially organic matter (humins) through leaves. The assimilation of all nutrients with this approach increases significantly, which in turn contributes to the development of plant specific substances and proteins that increase their resistance to disease, drought, high temperatures and the like.

Table 4 Grain yield of spring barley varieties depending on resource-saving optimization of nutrition, t/ha

| Variant | Number of treatments | Varieties (factor B) | | | | | | | | | |
|---|----------------------|----------------------|------|---------|-----------------------|------|-------------------|------|---------|-----------------------|------|
| | | Stalker | | | | | Vakula | | | | |
| | | years of research | | | the increase in yield | | years of research | | | the increase in yield | |
| | | 2016 | 2017 | average | t/ha | % | 2016 | 2017 | average | t/ha | % |
| Control | | 2.91 | 2.64 | 2.78 | 0 | 0 | 3.26 | 2.27 | 2.77 | 0 | 0 |
| 1. Fresh Floride 200 g/ha | 1 treatment | 3.67 | 2.75 | 3.21 | 0.43 | 15.5 | 4.07 | 2.43 | 3.25 | 0.48 | 17.3 |
| | 2 treatments | 3.92 | 3.08 | 3.5 | 0.72 | 25.9 | 4.38 | 2.92 | 3.65 | 0.88 | 31.8 |
| | 3 treatments | 4.04 | 3.34 | 3.69 | 0.91 | 32.7 | 4.52 | 3.24 | 3.88 | 1.11 | 40.1 |
| 1. Fresh Floride 300 g/ha | 1 treatment | 3.92 | 3.32 | 3.62 | 0.84 | 30.2 | 4.32 | 2.74 | 3.53 | 0.76 | 27.4 |
| | 2 treatments | 4.28 | 3.7 | 3.99 | 1.21 | 43.5 | 4.75 | 3.02 | 3.89 | 1.12 | 40.4 |
| | 3 treatments | 4.54 | 3.93 | 4.24 | 1.46 | 52.5 | 5.07 | 3.39 | 4.23 | 1.46 | 52.7 |
| 1. Fresh Floride 200 g/ha | 1 treatment | 3.69 | 2.74 | 3.22 | 0.44 | 15.8 | 4.13 | 2.29 | 3.21 | 0.44 | 15.9 |
| | 2 treatments | 3.9 | 3.09 | 3.5 | 0.72 | 25.9 | 4.33 | 2.45 | 3.39 | 0.62 | 22.4 |
| | 3 treatments | 3.97 | 3.37 | 3.67 | 0.89 | 32 | 4.45 | 2.59 | 3.52 | 0.75 | 27.1 |
| 1. D ₂ M ₁ 1000 g/ha | 1 treatment | 3.46 | 2.97 | 3.22 | 0.44 | 15.8 | 3.87 | 2.31 | 3.09 | 0.32 | 11.6 |
| | 2 treatments | 3.84 | 3.3 | 3.57 | 0.79 | 28.4 | 4.31 | 2.85 | 3.58 | 0.81 | 29.2 |
| | 3 treatments | 4.28 | 3.84 | 4.06 | 1.28 | 46 | 4.8 | 3.06 | 3.93 | 1.16 | 41.9 |
| 1. Escort 250 g/ha | 1 treatment | 3.48 | 2.7 | 3.09 | 0.31 | 11.2 | 3.86 | 2.36 | 3.11 | 0.34 | 12.3 |
| | 2 treatments | 3.82 | 3.03 | 3.43 | 0.65 | 23.4 | 4.25 | 2.75 | 3.5 | 0.73 | 26.4 |
| | 3 treatments | 4.15 | 3.39 | 3.77 | 0.99 | 35.6 | 4.65 | 3.05 | 3.85 | 1.08 | 39.0 |

In addition, nutrients, which are components of the growth-regulating substances, after contact with the leaf surface of the plants are rapidly absorbed, they pass the same way of synthesis as the elements that enter the plant through its root system, but 5 or more times faster. During seed treatment, biostimulators enhance the growth and

development of seedlings, in the subsequent phases of vegetation increase the intensity of photosynthetic activity of plants, in general, positively affect the yield and quality of grown products. Moreover, due to higher levels of crop yields due to use for power optimization growth-regulating substances, after the end of the growing season in the soil and remains much more plaintive-root residues, which in turn serve as organic fertilizer for the next crop. Thus, growth-regulating substances have a positive impact on the preservation of the main indicators of soil fertility.

For the southern Steppe zone of Ukraine, the presence of organic matter in the soil is extremely important for the accumulation, retention and further economical consumption of moisture by plants for crop formation. In compacted and depleted organic matter soil differences water in the rain and especially for high temperature conditions practically do not wet the soil, and evaporates quickly, that is, moisture is lost unproductively. The specified negative sign is not shown on fertile soils at high culture of agriculture. Consequently, even after prolonged drought, the moisture in these soils is retained and better used by plants. The importance of organic farming in this case increases, because in the South of Ukraine the main limiting factors for obtaining high yields of crops is the presence of moisture.

Our studies have determined that organic fertilizers in their properties can significantly improve the water-absorbing and water-retaining capacity of the soil, they act as mulch for the top layer of the soil, prevent evaporation of moisture and its unproductive losses. We show the above on the example of the influence of different types of organic fertilizers on the water-clay ability of the soil (Table 5).

Table 5 Water-absorbing ability of soil under the influence of organic fertilizers (average for three years)

| Experiment variant | Water absorbed, mm/h | % before control |
|-----------------------------|----------------------|------------------|
| Without fertilizer | 11,69 | 100,0 |
| Manure 30 t / ha | 13,60 | 116,0 |
| Green manure (peas) | 15,30 | 130,9 |
| Green fertilizer (rapeseed) | 14,73 | 126,0 |

Much more effectively moisture is used to optimize nutrition, which can be traced at the example of sunflower, which was grown without the introduction of organic and mineral fertilizers, and during the growing season only foliar feeding growth regulating substances were used.

For the conditions of the southern Steppe of Ukraine, which is characterized as a zone of unstable moisture and risky agriculture, when in some periods in the region for 100-120 days there is no precipitation, it is important to use moisture with the greatest efficiency – its reserves in the soil during the sowing of crops and precipitation that fell

during its growing season. We have determined the total water consumption of sunflower plants in the years of cultivation, its data are shown in table 6.

Table 6 Total water consumption of sunflower and its balance in the years of cultivation

| Years of research | Components of water consumption, m ³ / ha | | | Share of total water consumption, % | |
|---------------------------|--|--|---------------------------|-------------------------------------|--|
| | Soil moisture | Precipitation of the vegetation period | General water consumption | Soil moisture | Precipitation of the vegetation period |
| 2016 | 980 | 2293 | 3273 | 29.9 | 70.1 |
| 2017 | 670 | 1636 | 2306 | 29.1 | 70.9 |
| The average for 2016-2017 | 825 | 1965 | 2790 | 29.6 | 70.4 |

Significantly higher total water consumption was determined in a more favorable moist year – 2016, in which the soil moisture reserves were also large. However, in both years growing in the balance of total water consumption, the proportion of both soil moisture and precipitation of the growing season were almost the same. At the same time, it is most important to determine the coefficient of water consumption that characterizes the amount of moisture that the plant spends on the formation of the crop unit.

Sunflower crops on average for two years of research the least effectively used moisture without the use of biological products for foliar feeding. Thus, during treatment of crops with water only (control) plants spent 1320,4 m³/t, slightly less than the coefficient of water consumption already in processing plants in the phase of 3-4 pairs of leaves with retardin (0.25 kg/ha) 1211,9 m³/t. with Fresh energy depending on the doses of use – 1005.11-1171.6 m³/t, and these two drugs together: 941.7 is 1128 m³/t.

Carrying out fertilizing in the budding phase provided a more efficient use of moisture by sunflower plants. Among two feedings in the phase of 3-4 pairs of leaves budding and the coefficient of water consumption was reduced to 873.4-912.3 m³/t. Therefore only due to the processing of sunflower crops with growth regulators according to the background gopong application of mineral fertilizers N₁₆P₁₆K₁₆ it is possible to reduce the ratio of water consumption to 9.0-51.2 %.

It should be noted that since two years of sunflower cultivation, the water consumption coefficient has been determined somewhat higher in 2016, and in 2017, it was slightly less favorable in terms of precipitation.

The importance of biological products in improving the efficiency of water consumption of sunflower is illustrated in Fig. 3, where the better time for foliar application is clearly seen.

Use of moisture, m³/t




Figure 3. Influence of doses and timing of fertilizing with biopreparations on the coefficient of water consumption of sunflower (average for 2016-2017), m³/t

Treatment of plants with bio substances during budding is much more appropriate in comparison with the phase of formation of 3-4 pairs of leaves. Feeding in the budding phase according to the values of the water consumption coefficient slightly differs from their values in the processing of sowing twice in both phases of the development of sunflower plants.

CONCLUSION

Thus, as it is determined by the results of our research, the level of crop yields is significantly determined by the degree of soil fertility, namely it depends on its supply with the organic matter and the content of basic nutrients. It grows under the influence of fertilizers, the use of which on poorer soils contributes to a more significant increase in yield compared to fertile ones. A prominent place should be given to scientifically based alternation of crops in crop rotation, which improves the basic indicators of soil fertility, enriching it with organic matter and reduces the cost of production.

Due to the increase in organic matter in the soil, water absorption capacity and efficiency of moisture use by plants significantly increases. In the conditions of modern resource provision of the agricultural sector, in particular the high cost of mineral fertilizers, the missing volumes of organic application should be developed and modern approaches to plant nutrition should be implemented. The most effective and cheapest method is the use of biologics and growth regulating substances for presowing treatment of seeds or planting plants in the main periods of the growing season. This allows to reduce the dose of mineral fertilizers, increase crop yields, increase their resistance to drought and adverse environmental conditions, has a positive effect on the main indicators of quality of grown products, elements of fertility and significantly improves the efficiency of the use of plants moisture on the formation of the crop unit.

With such resource-saving approaches it is possible to achieve a certain stability in the management of agricultural crops, including in the zone of unstable moisture when changing soil and climatic conditions of the southern Steppe of Ukraine, which is the breadbasket, the source of vegetables, melons and other crops.

BIBLIOGRAPHICAL REFERENCES

1. Dobrovolskyi H. V. Soil conservation and fertility – the most important environmental problem of the XXI century // Soils and their fertility at the turn of the century. Materials of the II Congress of the Belarusian society of soil scientists. - Book 1. "Theoretical and applied problems of soil science". - Minsk, 2001. - p. 74-75.

2. Valentina Gamajunova Sustainability of Soil fertility in Southern Steppe of Ukraine, Depending on fertilizers and irrigation // Soil science Working for a Living Applications of Soil science to Present – Day Problems. - Springer International Publishing Switzerland 2017. – P. 159-166.

3. Medvedev V. V., Plisko I. V., Nakisko S. G., Titenko G. V. Soil Degradation in the world, the experience of its prevention and overcoming. Kharkiv: Stylish printing house, 2018. 168 p.

4. Baliuk S. A., Romaschenko M. I., Truskavetskyi R. S. Problems of ecological risks and prospects of land reclamation in Ukraine // Agrochemistry and soil science. Interdepartmental thematic scientific collection. Issue 87. Kharkiv: National scientific center named after A. N. Sokolovskyi", 2018. - p. 5-10.

5. Hamayunova V. V., Kovalenko O. A., Honenko L. H. Modern approaches to the management of the agricultural sector on the principles biologization and resource / Collective monograph "Rational use of resources in conditions of ecologically stable areas. Edited by P. V. Pysarenko, T. O. Chaika, I. O. Asmolov / Poltava, PDA, 2018.- p. 232-241.

6. Chaykovska L. O. Influence of Bio phosphor on plant productivity in conditions of southern steppe / L. O. Chaykovska // Optimization of structure of agricultural

landscapes and sustainable use of soil resources-K., 200. - p. 91-94.

7. Hamayunova V. V., Drobitko A.V., Panfilov A. V. Value of biodestruction of stubble for the biological agriculture in the conditions of southern steppe of Ukraine / / proceedings of abstracts of the international scientific-practical conference "Effective use of ecological agricultural production" - Scientific methodical center "Agrosvit". - K., November 2, 2017-p. 38-41.

8. Hamayunova V., Tuz M., Bazalii C. Impact of biologics on the features of consumption of legumes in conditions of southern steppe of Ukraine / Moldova, Stiinta Agricola, Nr.2,2017.- P. 23-29.

9. Filipiev I. D., Hamayunova V.V., Baliuk S. A., et al.. Systems of fertilizers of agricultural crops // Scientific bases of protection and rational use of irrigated lands of Ukraine / [Ed. By S. A. Baliuk, M. I. Romashchenko, V. A. Stashuk]. - K.: Agricultural science, 2009. - P. 279-299.

10. Miroshnichenko M. M., Hladkih Ye. Yu., Revtie-Uvarova A.V., et al. Nine approximations of the modern system of crop production // agricultural chemistry and soil science. Interdepartmental thematic scientific collection. Issue 87. Kharkiv: national scientific center named after A. N. Sokolovskyi", 2018. - p. 82-92.

Scientific development and achievements

Copyright © 2018 by Hamaiunova V., Hlushko T., Honenko L., Prylipko T., Bukalova N., Bogatko N., Shtuiak A., Dobrovolsky V., Stankevych S., Zabrodina I., Yevtushenko M., Biletskyj Y., Sychov M., Holubiev M., Voloshin N., Golovan L., Klymenko I., Chupryna Y., Lisovyy M., Fedorchuk S., Klymenko T., Trembitska O., Vasilyeva Ju., Lezhenina I., Kochmarskii V., Melnyk V., Skiba M., Vorobyova V., Vereschak V., Pasenko O., Bykov I., Vishnikin A., Al-Shwaiyat Mohammed E.A., Okovytyy S., Chernavskaya A., Hedjazi M., Zhuk T., Vorobyova V., Margarita S., Zabarna I., Prosaniiy S.

ALL RIGHTS RESERVED
Published by Sciemcee Publishing.
LP22772, 20-22 Wenlock Road
London, United Kingdom
N1 7GU

Sciemcee Publishing is part of SCIEMCEE.

It furthers the SCIEMCEE's mission by disseminating knowledge in the pursuit of education, learning and research at the highest international levels of excellence.

No part of this publication may be reproduced in any manner without the express written consent of the publisher, except in the case of brief excerpts in critical reviews or articles. All inquiries be address to Sciemcee Publishing, LP22772, 20-22 Wenlock Road, London, N1 7GU or publishing@sciemcee.com.

First Edition: 2018

A catalogue record for this publication is available from British Library.

Sciemcee Publishing has no responsibility for the persistence or accuracy of URLs for external or third-party internet referred in this publication, and does not guarantee that any content on such websites is, or will remain, accurate or appropriate.

Every effort has been made in preparing this book to provide accurate and up-to-date information which is in accord with accepted standards and practice at the time of publication. Nevertheless, the authors, editors and publishers can make no warranties that the information contained herein is totally free from error. The authors, editors and publishers therefore disclaim all liability for direct or consequential damages resulting from the use of material contained in this book. Readers are strongly advised to pay careful attention to information provided by the book.

Sciemcee Publishing also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

Sciemcee Publishing books may be purchased for educational, business, or sales promotional use. For information, please e-mail the Sciemcee Publishing at publishing@sciemcee.com.

ISBN 978-1-9993071-0-3

Includes bibliographical references and index.

